

This is a repository copy of *Childhood and adolescent influenza vaccination in Europe : A review of current policies and recommendations for the future*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/97790/>

Version: Accepted Version

---

**Article:**

McGuire, Alistair, Drummond, Mike [orcid.org/0000-0002-6126-0944](https://orcid.org/0000-0002-6126-0944) and Keeping, Sam (2016) Childhood and adolescent influenza vaccination in Europe : A review of current policies and recommendations for the future. *Expert Review of Vaccines*. pp. 659-670. ISSN 1744-8395

<https://doi.org/10.1586/14760584.2016.1138861>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



# Childhood and adolescent influenza vaccination in Europe: A review of current policies and recommendations for the future

Alistair McGuire, Mike Drummond & Sam Keeping

**To cite this article:** Alistair McGuire, Mike Drummond & Sam Keeping (2016): Childhood and adolescent influenza vaccination in Europe: A review of current policies and recommendations for the future, Expert Review of Vaccines

**To link to this article:** <http://dx.doi.org/10.1586/14760584.2016.1138861>



Accepted author version posted online: 06 Jan 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

**Publisher:** Taylor & Francis

**Journal:** *Expert Review of Vaccines*

**DOI:** 10.1586/14760584.2016.1138861

**Childhood and adolescent influenza vaccination in Europe:  
A review of current policies and recommendations for the  
future**

**Alistair McGuire<sup>a</sup>, Mike Drummond<sup>b</sup>, Sam Keeping<sup>a</sup>**

<sup>a</sup> *LSE Health, LSE, Houghton Street, London WC2 2AE.*

<sup>b</sup> *Centre for Health Economics; Alcuin 'A' Block; University of York; Heslington, York;  
YO10 5DD, UK*

**Correspondence:**

Professor Alistair McGuire,  
LSE Health  
London School of Economics  
Houghton Street  
London  
WC2A 2AE

+44 (0) 207 955 6375

[A.J.MCGUIRE@LSE.AC.UK](mailto:A.J.MCGUIRE@LSE.AC.UK)

## Summary

Children and adolescents experience some of the highest rates of influenza infection and the subsequent burden on both infected children and their parents/carers is substantial. Vaccinating children and adolescents against seasonal influenza has the potential to reduce the burden of disease in both vaccinated and unvaccinated individuals due to the pivotal role that younger age groups play in the transmission of infection. While countries such as the USA, Canada and the UK have consequently recommended the universal vaccination of children, the vast majority of European countries have not yet extended their vaccination policies to this age group. This review examines the rationale for childhood and adolescent vaccination against seasonal influenza and reviews current vaccination policies in Europe. We discuss key policy considerations for European countries that must be considered when extending vaccination programmes to younger age groups alongside recommendations for European policy makers based on our findings.

**Keywords:** Influenza, vaccine, childhood, adolescent, policy, Europe

## Introduction

Influenza is a highly infectious virus that affects the respiratory tract. Influenza viruses are categorized as either type A, B or C, with the majority of clinical illness caused by types A and B [1].

Although otherwise healthy individuals who contract the virus can usually expect to make a full recovery, the virus may have serious consequences for some groups, such as the very young, the elderly and those with chronic illnesses [2]. Its impact on morbidity and mortality amongst these groups is significant; The European Centre for Disease Prevention and Control (ECDC) estimates that each year the average number of people in Europe who die prematurely due to influenza infection is 40,000 [3]. The virus also places significant strain on health systems when its circulation is at its peak [4].

Countries that have employed mass influenza vaccination have historically targeted those most at risk of serious complications. However, in the last 10 years some health authorities have also opted to vaccinate healthy children. Children experience some of the highest attack rates during influenza outbreaks [5] and also play a key role in the spread of the virus [6], although they do not experience the same rates of complications or death as at risk-groups [7]. Vaccinating children has therefore been seen by some as an effective way to deliver health gains through the direct protection of children themselves and also indirect protection of others, such as relatives and family members, who may have otherwise come into contact with the virus. However, the inclusion of healthy children and adolescents within seasonal influenza vaccination programmes remains far less widespread than for the elderly or people with predisposing conditions.

This review examines the rationale for childhood and adolescent vaccination against seasonal influenza and reviews current vaccination policies in European countries. We discuss key policy considerations for European countries that need to be considered when extending vaccination programmes to younger age groups and provide recommendations for European policy makers based on our findings.

# **Rationale for the seasonal vaccination of children and adolescents against influenza**

For those countries with current influenza vaccination programmes for children and adolescents, the detailed reasons underpinning vaccination policy differ between them but can nonetheless be grouped into two broad categories: (1) reducing the burden of disease in younger age groups through direct protection, and (2) indirectly protecting high-risk groups from disease by disrupting community transmission (also known as “herd protection”).

## **Direct protection**

The burden of influenza amongst children is substantial, with attack rates reaching 20–30% of the population on average in each influenza season [8]. There is the potential for more serious complications in children such as acute otitis media [9], bacterial co-infections such as pneumonia [8] or methicillin-resistant *Staphylococcus aureus* [10], and febrile seizures [11]. Vaccination against influenza has been shown to reduce the incidence of acute otitis media by over 35% [12,13] and reduce the chance of contracting more serious complications such as pneumonia [14]. There have been isolated reports that uncomplicated and complicated influenza or influenza-like illness (ILI) in children and adolescents has a significant impact on health-related quality of life [15]. Influenza cases among children and adolescents also place a substantial burden on healthcare systems in terms of resource utilisation. Data from the USA during the 2002–2003 and 2003–2004 influenza seasons, for example, showed that medical clinic and emergency department visits due to laboratory-confirmed influenza in children <5 years were estimated to be in the range of 50–95 and 6–27 per 1000 children, respectively [16]. The risk of

hospitalization for healthy children generally decreases with age, with children aged  $\leq 4$  years experiencing rates similar to other groups, who have traditionally been targeted for vaccination [7].

A number of estimates are available for the indirect costs of influenza in adults due to both absence from work and reduced productivity while at work (known as presenteeism) [17,18]. The specific impact that childhood influenza has on the wider economy due to parents and responsible individuals taking time off work to care for those who become sick has also been investigated. A prospective cohort study carried out in three large US companies found that employees with at least one child in their household with self-reported acute respiratory tract infection (ARI) reported significantly higher levels of absenteeism due to household illness or injury (mean 0.8 vs 0.3 days per month) than those with no children reporting ARI [19]. Those with children who had ARI that met the criteria for ILI also had greater lengths of absence due to household illness or injury than others (0.9 vs 0.5 days per month). Although these results should be treated with caution as they rely on self-reported illness, they do give some indication of the extent to which influenza in children impacts those around them.

The above evidence on the negative impact of influenza in children, along with the availability of safe and effective vaccines and a desire to increase uptake in children at risk of complications, provided the rationale for the USA's Advisory Committee on Immunization's (ACIP) decision to recommend all children aged 6 months and over to be vaccinated against influenza on an annual basis [20]. A similar argument was put forward by the Canadian National Advisory Committee on Immunisation (NACI) in recommending the vaccination of children aged 6 months to



<2 years in the 2011–2012 influenza season [21], with the upper age limit extended to <5 years in the following season [22].

Two recent systematic reviews on the burden of childhood and adolescent influenza in Europe reveal a similar picture to the one seen in North America, with high rates of infection and healthcare utilisation [23,24]. For example, data drawn from England [7] over the course of eight consecutive seasons (2000–2001 to 2007–2008) show similar rates of general practitioner consultations (61 per 1000) and hospitalizations (two per 1000) for children aged 6 months to 4 years to those seen in the USA. Estimates of the proportion of parents having to take time off work to care for a child with laboratory-confirmed influenza range between 11.2 and 61% for Western Europe [23]. The average number of days absent ranged from 1.3 to 6.3 (median range 2–4) per influenza case [23]. A study in Sweden found clear co-variation of seasonal ILI visits and temporary parental absence from work due to caring for ill children [25]. Parental absenteeism due to childhood influenza is expected to increase as female labour market participation for those aged between 20 and 64 years moves towards the 75% target set by the European Commission in 2010 [26].

### **Indirect protection**

The rate at which infections emerge within a population depends largely on the frequency and type of contact which occurs between individuals, the residual level of immunity within the population (i.e., the size of the susceptible population), the effectiveness of available vaccines [27] and the basic reproductive number,  $R_0$ .  $R_0$  is defined as the average number of secondary infections due to a single case when placed in an entirely susceptible population [28];  $R_0 \geq 1$  is required for a virus to

remain in circulation. The  $R_0$  for influenza has historically ranged anywhere from 1.1 to 3.7 [28]. The reproduction number can be used to determine the threshold for the proportion of protected individuals above which transmission will be disrupted ( $1 - [1/R_0]$ ), with higher reproduction values requiring greater levels of protection.

Reaching this threshold for influenza is challenging for a number of reasons. Firstly, the continually evolving nature of influenza viruses means that the degree of susceptibility within populations remains high from year to year, particularly for type A viruses [29]. In addition, older children and young adults are particularly susceptible to influenza B [30], the incidence of which can vary dramatically between seasons [31]. Some mutations can also make the virus more infectious, which can in turn increase the value of  $R_0$  [29]. Secondly, available vaccines are not 100% effective, for example in seasons where there is a poor match between the vaccine strains and the circulating virus and for populations whose immune systems are weaker, such as the elderly [19]. As the critical threshold for protection is proportional to the effectiveness of vaccination, low vaccine effectiveness can mean prohibitively high values for the critical threshold for protection [28].

The increased frequency of contact amongst children as well as between them and adults, relative to other age groups [6], gives rise to the possibility of targeting children and adolescents for influenza vaccination to provide indirect protection to other members of the population through reduced transmission [32], as well as direct protection to the children. The generally higher vaccine efficacy of newer influenza vaccines (e.g., live attenuated, already licensed and available, [33] and adjuvanted influenza vaccines, yet to be licensed in most countries for children [34], means the level of coverage in this population required to reduce incidence

among other age groups may be lower than for groups more commonly targeted for vaccination, such as the elderly. The low effectiveness of live attenuated influenza vaccine (LAIV) noted in the 2013–2014 season in the USA is believed to be a result of heat exposure during shipping and the poor thermal stability of A/California H1N1 LAIV strain from a pre-existing mutation in the stalk region in the master virus seed [35]. A more stable strain was selected for the 2015–2016 season [35].

Data derived from a nu

mber of observational studies demonstrate the impact that vaccinating younger age groups can have on influenza transmission [36–38]. Japan was the first country to implement routine vaccination of healthy schoolchildren after experts identified them as key spreaders of infection during a series of major epidemics during the 1950s [39]. Between 1962 and 1987 when the routine programme was in operation, mortality attributable to influenza and pneumonia fell by 10,000–12,000 deaths per year, while all cause-mortality declined by 37,000–49,000 deaths per year. Importantly, no comparable decline was observed in the USA over the same period (where no childhood influenza vaccination programme was in place) and deaths in Japan promptly rose after the programme was discontinued, which when taken together appear to suggest that the decline in mortality was strongly associated with the childhood influenza vaccination. However, it should be noted that the prevalence of multi-generational homes in Japan at the time may have contributed to the herd protective effect, as family members were more likely to be exposed to influenza infection from close relatives (particularly children).

As many observational studies of the effect of influenza vaccines lack laboratory-confirmed viral endpoints [40], there has been a desire for randomized

clinical trials (RCTs) to confirm the presence of indirect protection. One classic example is a cluster randomized trial of trivalent inactivated influenza vaccine (TIV) delivered to children aged between 36 months and 15 years in Hutterite communities in western Canada. The study showed that the vaccine was 61% effective at preventing influenza among unvaccinated persons in communities randomized to receive the vaccine [41]. A similar interventional study of the effectiveness of school-based influenza vaccination found that most outcomes related to ILI were significantly lower in households where the child had received vaccination compared to control households where the child had not [42]. The results of studies such as these, which demonstrate indirect protection from childhood and adolescent vaccination against influenza, have constituted the evidence base used to recommend universal influenza vaccination for those aged  $\geq 6$  months in the USA [43].

## Overview of existing programmes

For the purposes of reviewing current vaccination policy towards influenza among all countries in the European Union, we have placed them into six regions based on geographical proximity. The groups and their constituent countries can be found in TABLE 1.

Only eight countries currently have a general recommendation for the vaccination of children and/or adolescents against influenza: Austria, Finland, Latvia, Malta, Poland, Slovakia, Slovenia and the UK (FIGURE 1). Of these, only Finland, Latvia and the UK provide the vaccine free of charge. FIGURE 2 shows how these countries are distributed among the different regions and indicates that no single region favours the strategy of vaccinating children and adolescents over another. A brief overview of the existing recommendations in each region and their levels of vaccine coverage is provided below, with data drawn from the ECDC [44] and the Vaccine European New Integrated Collaborative Effort (VENICE) survey of countries for the 2011–12 influenza season [45].

### Baltic countries

Latvia is the only Baltic country with a positive vaccination recommendation for the young. Health authorities provide TIV free of charge to all children aged  $\geq 6$  months and  $< 3$  years, as well as to those aged  $\geq 65$  years. This is in keeping with its general approach to vaccination policy, which is amongst the most aggressive in Europe [46]. For example, vaccination is mandatory for all state employees and Latvian health care providers, with the latter required to obtain a signature from any person who refuses to be vaccinated as evidence that all relevant information on the risk of disease has been provided. Despite these measures vaccination coverage remains

very low according to the most recent VENICE group survey (0.1% for those aged 6–23 months) [45].

### **Central European countries**

Both Poland and Austria recommend the vaccination of children and adolescents using TIV, the latter opting to vaccinate from  $\geq 7$  months to  $< 16$  years and the former from  $\geq 13$  months to  $< 18$  years. The recommendations in both countries were originally made in 2010 [47] although Austria subsequently reduced the upper limit from  $< 18$  to  $< 16$  years [48]. Neither country provides public funding for these programmes nor has a high vaccination coverage rate (2.4% for those aged 5–14 years). Germany and Liechtenstein both provide influenza vaccine free of charge for persons aged  $\geq 60$  and  $\geq 65$  years, respectively, with the German state of Saxony including those aged  $\geq 6$  months within their recommendations [49].

### **Eastern European countries**

In Eastern Europe, Slovenia and Slovakia recommend seasonal vaccination of children, the latter for ages  $\geq 6$  months to  $< 13$  years and the former for ages  $\geq 6$  months to  $< 3$  years, with no reimbursement of vaccine costs. Vaccination rates are below 3% in both countries [45]. Slovenian vaccination policy differs from most other European countries in that childhood vaccination across nine disease areas is mandatory (not including influenza), with no exceptions based on religious belief and fines for those who fail to comply [46]. The remaining countries do not have childhood or adolescent vaccination policies, but have policies for adults [44].

### **Nordic countries**

Finland has had a publicly funded vaccination programme for all healthy children aged 6–35 months since 2007–2008 [50]. Importantly, this is part of their routine

schedule and as a result they have achieved higher levels of uptake than other countries with a childhood recommendation (36.2% in 2007–2008) [51]. Iceland also provides reimbursement for vaccine but only for those aged  $\geq 60$  years.

### **Southern European countries**

Among Southern European countries only Malta has a positive recommendation for childhood influenza vaccination ( $\geq 6$  months to  $< 3$  years) but this is not part of the routine schedule.

### **Western European countries**

The UK government's advisory body on vaccine policy, the Joint Committee on Vaccination and Immunisation (JCVI), issued a recommendation that all healthy children and adolescents aged  $\geq 2$  to  $< 17$  years be vaccinated using LAIV [52]. The intention is for the programme to be phased in over a number of years, starting in the 2013–2014 season. The pilot programmes introduced in the 2013–2014 and 2014–2015 seasons were mainly delivered through schools and achieved promising results, with an average uptake of 52.5% in 2013–2014 and 56.8% in 2014–2015 in primary schools in England [53,54]. Furthermore, results from the 2014–2015 season revealed that vaccinating primary school age children in pilot areas resulted in significant and non-significant reductions in a range of surveillance indicators for both the targeted age groups and wider society (due to herd protection) [54]. No other Western European countries currently have recommendations for childhood or adolescent vaccination.

## **Key policy considerations for the introduction of childhood and adolescent vaccination programmes in Europe**

Our review reveals that fewer than half of European countries currently have recommendations for healthy childhood and adolescent vaccination (FIGURE 2), despite evidence to support the vaccination of this age group as a strategy to reduce the burden of influenza. The limited roll out of influenza vaccination for healthy children, along with the variation in recommendations among countries who have them, is confirmation that a European consensus has yet to be reached over the policy.

Where data are available, they indicate that vaccination rates in age groups with a national recommendation are very low. In addition, only three countries, Latvia, Finland and the UK, have public funding for childhood and adolescent vaccination programmes. Even with public funding, however, vaccination rates are extremely low in Latvia, especially when compared with the good uptake achieved in the first year of the UK's childhood vaccination programme [45,53]. This suggests that a complex set of factors contribute to the success of childhood and adolescent vaccination programmes and need be considered by policy makers. Below, we consider key policy considerations for European countries considering the implementation of childhood and adolescent influenza vaccination programmes and summarise our recommendations.

### **Evidence of vaccine efficacy and safety**

The starting point for policy makers when evaluating whether a new vaccine should be recommended and introduced into the schedule is evidence of efficacy and safety. While evidence is usually provided through a RCT, the need for the rapid



evaluation of influenza vaccines due to changing compositions from season to season and infeasibility of carrying out annual RCTs has led the European Medicines Agency to develop a unique set of processes for their authorization for human use, which in some circumstances permits the use of data from comparative studies on immunogenicity instead of that which would normally come from an RCT [55].

Nevertheless, a number of different meta-analyses of RCTs have examined the efficacy of inactivated and LAIVs administered to healthy children. These studies have demonstrated good efficacy for LAIV in children aged >2 years [56,57] and superior efficacy compared with TIV [56]. In contrast, there is limited evidence for the efficacy of influenza vaccines in children aged <2 years, the group most at risk of complications and death due to the virus. As for adjuvanted vaccines, a phase III trial carried out in 2011 found it to provide 83% immunological efficacy (95% CI 74–93%) in children aged from 6 months to <6 years, compared to just 43% (95% CI 15–61%) for standard TIV [58]. However, this vaccine is currently unlicensed in Europe [59]. Countries will have to consider the varying efficacy between age groups when selecting target populations for new vaccination programmes and any safety signals, such as the recent signal for narcolepsy seen with the pandemic influenza vaccine [60].

A number of observational studies have been carried out in European children to determine the effectiveness of influenza vaccines in real world conditions. An observational cohort study carried out in Finland showed TIV to be 83% (95% CI 58–93%) effective at preventing influenza in children aged 7–50 months [12]. Another similar Finnish study of TIV showed effectiveness against influenza A to be 84%

(95% CI 40–96%) and 45% (95% CI –34 to 78%) for influenza B [61]. A multi-centre case-control study of inactivated influenza vaccine effectiveness during the 2012–2013 season found that, after adjusting for the impact of comorbidities, the pooled vaccine effectiveness across inactivated influenza vaccine types in those aged 0–14 years was 22% (95% CI –37 to 56%), 37% (95% CI –44 to 72%) and 36% (95% CI –41 to 71%) against influenza B, influenza A (H1N1) and influenza A (H3N2), respectively [62].

There are currently limited effectiveness data for live attenuated and adjuvanted vaccines in European populations due to their limited usage in routine practice. Results from the 2014–2015 season in the UK, which was characterised by the circulation of drifted influenza A and B viruses, found that LAIV provided non-significant positive protection against influenza A and significant protection against B [63]. These results support literature that indicates that LAIV can provide cross-protection against drifted strains [63].

The apparent lack of data on effectiveness of both TIV and LAIV in healthy children has been cited as a significant barrier to the wider implementation of childhood influenza vaccination [24,64]. However, this is largely due to the vaccines being used sparingly in this age group across Europe. The UK programme is likely to fill this gap for LAIV in years to come, yet the willingness of health authorities in the UK to introduce the vaccine based on current evidence and modelling suggests that in the meantime other countries may choose to implement their own programmes rather than wait for this information to become available.

## Local evidence to support the impact of vaccination

While further RCTs such as the one carried out Loeb *et al* [41] would be of great assistance to other countries considering whether or not to introduce the routine vaccination of children and adolescents against influenza, the scope to conduct such trials in the future is heavily constrained on the grounds of practicality (randomizing entire communities is unfeasible), cost, and whether such studies are ethical [40]. In the absence of RCT evidence, observation studies with virologically confirmed endpoints are widely regarded as the next best option for clinical studies of influenza vaccination, but also require resources which may not be readily available in all countries [40]. A solution to this problem is to utilize mathematical modelling to produce estimates for the likely direct and indirect impact of vaccination programmes on seasonal influenza epidemics in terms of health and economic outcomes [27]. This approach is appealing as models are relatively cheap to produce compared to large trials, although the data requirements, such as virological surveillance and contact pattern data, can be burdensome to collect if not already available [65]. Another limitation of such models is that they require various assumptions to be made, such as levels of pre-existing immunity within the population, which makes their predictions uncertain. This can be mitigated by carrying out sensitivity analyses to test the robustness of model findings.

The UK's JCVI has a long history of using mathematical models to inform its policy recommendations. For example, the UK was the first country to implement mass vaccination against meningitis C, with the decision largely based on the predictions of a dynamic transmission model [66]. Modelling work done by the Health Protection Agency, now Public Health England, who support the work of the JCVI, showed that the impact of extending the seasonal vaccination programme in the UK

to include those aged from 2 to 18 years (using LAIV) was likely to be highly cost-effective based on the UK threshold [52]. A similar analysis published around the same time noted that vaccinating this age group was estimated to result in net cost savings among unvaccinated members of the population, with the greatest impact seen among the elderly even though they themselves had high pre-existing vaccination coverage (75%) [67]. In 2012, the JCVI considered the evidence and recommended a staggered roll out of an extended vaccination programme for those aged 2–16 years. The rationale for the phased introductions was that it would allow for additional capacity to deliver the vaccines in schools to be created, while at the same time limiting the disruption to existing services. Subsequent modelling work has gone even further and suggested that targeting infants and adolescents rather than at risk groups at the time when the programme was first implemented may have been a more efficient strategy for reducing the burden of influenza [68]. The phased roll out of the UK's programme has allowed for early assessment of its effects, with early evidence suggesting indirect protection may be occurring [53]. Such an approach may be appealing to other countries as it avoids inertia from the continuing absence of direct evidence while mitigating the risk of required capital investments. A similar cost-effectiveness analysis from Finland concluded that influenza vaccination would be cost saving in all children  $\leq 13$  years [69].

### **Healthcare system factors**

The structure of a country's healthcare system places a number of constraints on decision makers when determining optimal vaccine policy [70]. Important considerations include whether policies are made at the national or local level [70], the availability of funding and reimbursement for vaccines and existing capacity to deliver vaccination programmes [71], for example through schools. There are also

interdependencies between these areas which influence the feasibility of implementing childhood influenza vaccination. Many countries have created National Immunisation Technical Advisory Groups (NITAGs), which sit either within government or independent of it, to produce guidance on changes to the existing schedule of vaccinations as well as the introduction of new vaccines [72]. However, there is heterogeneity across NITAGs in both their capabilities and the extent to which their recommendations impose a legal requirement on governments to ensure their implementation [73].

In England, the Health Protection (Vaccination) Regulations 2009 requires “so far as is reasonably practicable, that the recommendations of JCVI are implemented”, so long as their assessments demonstrate cost-effectiveness [74]. The JCVI is also supported by staff from Public Health England, the national agency responsible for health protection and promotion, which has access to high quality virologic surveillance data along with extensive information on resource use [68]. This is made possible by the highly centralized nature of the National Health Service in England, along with the long-term investment that has been made in the infrastructure to allow both epidemiological data collection and analysis [75]. Centralized systems can also take advantage of their monopsony powers to negotiate lower vaccine prices [76]. This has been the case in England as well as the other UK territories, where a centralized system has also helped foster a system of school nurses who are able to facilitate mass vaccination of children in schools [77].

The pre-existing capacity to deliver vaccination through schools is important for a number of reasons. Firstly, a minimum level of coverage must be achieved among children and adolescents in order for community transmission to be disrupted, and

consequently for the benefits from protection to be realised [28]. Although some progress has been made in increasing uptake in countries which already have these programmes such as Finland and the USA, uptake remains suboptimal [78]. School-based mass vaccination programmes can potentially deliver higher levels of uptake and they have been used successfully in the USA for a number of years [42], although robust data of their actual effect is limited [79]. The first year of the UK's extended influenza vaccination programme for children carried out a number of school-based pilots, which suggested that a variety of school settings are feasible and can achieve a good level of coverage [80]. School-based programmes have the added benefit that the vaccine can be delivered prior to onset of the influenza season and can also serve to reduce the administration costs of delivering influenza vaccines, which have been recently shown to be slightly higher for LAIV than TIV [81]. However, there are barriers to implementing school-based systems: creating or enhancing the healthcare infrastructure along with educational provider arrangements can be challenging and costly.

Nevertheless, previous experience has demonstrated that childhood vaccination programmes can be delivered through private (e.g., USA) [82] or public systems (e.g., UK) [53] with good uptake and through a number of different services such as primary healthcare providers, nurses, healthcare support workers and school nurses [80]. Countries looking to implement such programmes will likely have to adapt programmes to their existing infrastructure or alternatively invest in developing healthcare infrastructure.

## Economic considerations

Providing vaccines free of charge has also been shown to have an important effect on uptake [83,84]. In countries where the assessment process for vaccines contains a health economic component, the cost-effectiveness of new programmes, such as influenza vaccination for healthy children, is likely to be highly sensitive to both the costs of the vaccine and its administration. In these circumstances, the ability of those responsible for procuring the vaccine to negotiate a favourable price, either through volume discounts or tender processes, becomes a key determinant of whether new vaccines can be introduced. In contrast, systems with no central funding may be more able to deliver recommendations which are broader in scope due to the absence of a budget constraint. A counterbalance to this is that if the vaccine is to be funded via insurance programmes or out of pocket expenditures, the absence of national risk pooling reduces the value of vaccination to the individual. This is because the indirect benefits from herd protection are less important in individual consumption decisions, possibly leading to lower coverage. In this case, policy makers might focus their own limited resources on exploring the viability of programmes which are viewed as having a greater likelihood of success.

Reimbursement of vaccine alone may not be able to guarantee sufficient levels of coverage to justify vaccinating children against influenza, however. Another major driver of vaccine coverage is physician engagement with the programme and the perceived benefit, both to themselves and their patients [85]. While centrally or regionally procured, vaccinations can improve the cost-effectiveness profile of vaccination programmes at the population level, they provide weak financial incentives to private physicians as time spent administering the vaccine could be substituted for more lucrative activities. This is an important issue in the context of an

intervention that has low perceived health benefit, such as influenza prevention in the pediatric population. As a result of this, many countries offer financial incentives to physicians to ensure sufficient levels of coverage can be reached [84]. While research on the effectiveness of such incentives is generally of a low quality, it does support the use of such incentives [86]. These create an additional cost for vaccination programmes which become part of the implementation decision.

TABLE 2 gives an overview of health system and vaccination policy features for a sample of European countries [72]. Countries like the UK, Ireland and Sweden who have tax funded healthcare systems with national risk pooling, tender procurement processes and the infrastructure for delivering vaccinations through both primary care and schools appear to be those most suited to implementing seasonal influenza vaccines. At the other end of the spectrum are countries like Germany, France and Belgium where vaccinations are primarily paid for by social health insurance schemes, funded through employee and employer payroll contributions, and largely administered by private physicians. It is therefore perhaps understandable that these countries have yet to move to a more comprehensive influenza vaccination programme for healthy children and adolescents.

### **Demographic factors**

Decisions regarding the best way to combat the threat posed by seasonal influenza epidemics are heavily influenced by the characteristics of the populations which are at risk, as these are key drivers of disease epidemiology. Clearly the size of the school-aged population is an important determinant of the impact that influenza cases have on both child and parental wellbeing as well the wider health system and economy. FIGURE 3 shows the proportion of the population aged 0–14 years for



selected European countries. Of the countries with a recommendation for childhood vaccination (those highlighted in red), only the UK and Finland have proportions of school-aged children above the median of 15.4%. However, a relatively large young population does not guarantee that the burden of influenza among them will be significant, with variables that influence this also having important implications for whether vaccinating children can provide indirect protection to at risk groups. A distinction should be drawn between high-risk groups (e.g., the elderly, infants and those with predisposing health conditions) and high transmission groups (e.g., school-aged children) [87]. In some circumstance both groups may overlap but usually the amount of transmission from child-to-child and from children to other age groups are key determinants of overall burden, and to a large degree depend on demographic variables such as household size, levels of urbanization and social contact patterns [6].

The profile of national workforces is another important factor when considering the potential benefit of vaccinating children against influenza. Female labour market participation has been steadily increasing since the early 1990s [88] and is expected to increase in the coming years. Therefore, the proportion of children for whom both parents are in paid employment is also likely to rise. This limited excess capacity in labour markets makes protecting children against influenza much more appealing than in circumstances where staff off sick can easily be replaced. Furthermore, if older relatives are expected to act as informal care providers under these circumstances, this could increase transmission between children and those most at risk of complications, which further strengthens the case for vaccinating children in the first place.

## Cultural factors

The ethical and practical justification for exposing otherwise healthy young people to the risks associated with the vaccine, no matter how small, to protect primarily themselves, but also others, depends largely on the prevailing cultural trends in the countries in question and has to be taken in to account when discussing vaccination policy. It could be argued that countries with existing vaccine schedules that achieve high coverage already demonstrate that the social value of vaccination is well understood by the public, as the individual risk trade-off in the case of diseases with extremely low incidence such as diphtheria and polio may not be sufficient alone to justify their continuing acceptability among parents. However, the inclusion of these diseases within combination vaccinations, which distributes the disutility of adverse events across diseases, may partly explain why uptake has not fallen.

From an ethical standpoint, the acceptability of the vaccine in question by different faith groups also has to be considered. In the UK, the use of porcine gelatine in the LAIV used in the childhood immunization programme drew attention in the media. However, many multi-faith groups have deemed the vaccines acceptable and Public Health England has published advice from representatives of faith communities [89].

An important practical consideration is the acceptance of the vaccine from the general public and the sensitivity of public opinion regarding external messaging on the safety and efficacy of the vaccine [90], particularly in light of recent anti-vaccination movements. Latvia had the highest proportion of respondents (82%) who had no intention to take the seasonal influenza vaccine in a recent Eurobarometer survey, suggesting that strong scepticism among the population about the need for

vaccination could be driving the low influenza vaccination rates despite a publicly-funded programme.

The agency model of medicine in which patients handover responsibility for care decisions can become problematic when the views of perceived sources of expertise, such as media outlets, are not supported by empirical evidence (for example, as happened with the measles, mumps and rubella vaccine controversy in the UK [91]). As the downstream effects of falls in coverage can be highly detrimental for public health [92], providing accurate, clear, information regarding the safety and/or effectiveness of vaccination is essential to gain public confidence. For example, the JCVI have explicitly stated that in rolling out the expanded influenza vaccination programme in the UK, extra care must be taken to “inform and educate parents, children, healthcare professionals and others about influenza, the live attenuated intranasal vaccine and the benefits of the extending the programme to children and to the wider population...” [52]. Attitudinal research in England and Scotland revealed that, in general, the introduction of the expanded vaccination programme with LAIV was well-accepted by children and their parents [80]. While the UK has had success in expanding its schedule while maintaining high levels of uptake, it remains to be seen whether this will continue with the roll out to older age groups and, if so, whether other European countries attempt to replicate the approach.

Overall, it is clear that childhood and adolescent vaccination against influenza can offer substantial benefits, although these will differ in scale between different countries. Here, we identified a number of issues that need to be considered by

countries looking to implement a new childhood and adolescent influenza vaccination programme. We provide a summary of our recommendations below.

## **Recommendations**

- Ensure robust efficacy and safety data exist for the vaccine in question.
- Utilize modelling to provide country- or region-specific data on the health or economic impact of vaccination to support policy and payer decision making.
- Consider existing healthcare infrastructure that can be utilized for a mass vaccination programme and the logistics and expense of developing new infrastructure.
- Utilize experiences from other countries to guide the implementation of local programmes.
- Consider local demographics that may affect the potential impact of the vaccine.
- Develop country-specific solutions to enable to sufficient uptake of the vaccine for the levels of indirect and direct protection desired, e.g., reimbursement for all or some of the target population and physician engagement.
- Develop clear and consistent messaging for the public surrounding the efficacy and safety of the vaccine.
- Anticipate local cultural issues that may affect the reception of the vaccine and determine strategies to manage any issues.

## Conclusions

The unpredictability of the influenza virus continues to present a major challenge to healthcare professionals and policy makers alike. Vaccination nonetheless remains the most effective means of reducing the incidence and severity of influenza, yet uptake of the vaccine in many European countries remains suboptimal. The increased use of childhood vaccination is an opportunity to not only reduce the substantial burden of disease in this age group, it can potentially also help to close gaps in protection for those most at risk of serious complications due to the pivotal role they play in the spread of the virus.

Major barriers to the addition of influenza vaccination to routine childhood schedules include questions over vaccine effectiveness, especially in those aged <2 years, its impact on transmission, acceptability among parents and guardians, and the costs of implementation. While many countries will be closely watching the roll out of the UK programme for answers to these questions, and ultimately to the question of whether they should implement their own, choosing to set up their own limited pilot programmes, guided by existing data, may lead to better outcomes in the long run as these can be properly tailored to local circumstances. Countries with younger populations, a high disease burden and population density along with low levels of coverage and the possibility of delivering the vaccine within their existing architecture should be prime candidates.

## Expert commentary

It is becoming increasingly apparent that children bear a high burden of influenza, which has historically been underestimated [23]. A growing body of evidence indicates that the vaccination of children can reduce the burden of influenza in children themselves, and also provide indirect “herd protection” benefits to the wider community [41,42]. The implementation of childhood and adolescent vaccination programmes is therefore a strategic approach to control influenza infections and reduce the burden of disease in the population. The UK’s Joint Committee on Vaccination and Immunisation [52] agreed with this view, and in 2012 issued a recommendation that the UK implement a childhood immunization programme, starting in the 2013–2014 influenza season. Preliminary results from the UK’s childhood vaccination programme against influenza indicate that a national programme delivered to children either through general practitioners or school-based programmes can achieve a good uptake and may provide indirect benefits to the wider population [53,80]. However, other European countries have seemed reluctant to include healthy children and adolescents within seasonal influenza vaccination programmes, with only a minority currently having such programmes. The current situation highlights a European consensus has yet to be reached over the policy. Countries may be waiting for further external data on the impact of childhood and adolescent vaccination programmes, but we believe that there is sufficient evidence for European countries to introduce pilot programmes for the vaccination of healthy children against influenza. Local pilot programmes will enable countries to gather data to tailor their programmes to local circumstances and support future national programmes. We hope that the policy considerations discussed in this article will

provide countries with guidance to implement national childhood and adolescent vaccination programmes for influenza.

## Five-year view

Childhood and adolescent influenza vaccination programmes are likely to become increasingly common across Europe in the future, despite some initial hesitance to their adoption. Countries with existing childhood and adolescent vaccination programmes – and public funding for vaccines – are ideal candidates for the introduction of a childhood and adolescent influenza vaccination programme and are likely to be the first to introduce such programmes. Countries considering the implementation of childhood and adolescent vaccination programmes are likely to look to the UK in the coming years as the UK extends its national childhood vaccination programme to older age groups. By 2018–2019, the UK is expected to have rolled out the programme to children aged 2–16 years [52,93]. Data from successive seasons of the UK's programme will be eagerly awaited by countries looking to assess the impact of childhood and adolescent vaccination programmes in Europe. However, countries will also have to consider local and regional factors that may affect the implementation of such programmes.



## Key issues

- Children and adolescents experience some of the highest rates of influenza infection during seasonal epidemics. The negative effects of influenza on both infected children and their parents/carers is substantial.
- Vaccinating children against seasonal influenza has the potential to reduce the burden of disease in both vaccinated and unvaccinated individuals due to the pivotal role that younger age groups play in the transmission of infection.
- Vaccination programmes may have additional benefits, including reductions in absenteeism resulting from parents/carers taking time off work to care for sick children and reductions in pressure on healthcare services during peak influenza circulation.
- Large developed countries such as the USA, Canada and the UK now recommend influenza vaccination of healthy children and many have school-based influenza vaccination programmes.
- However, the vast majority of European countries have chosen not to implement their own versions of these programmes, and only a minority of countries with recommendations have public funding for influenza vaccines. Possible explanations for this include continuing uncertainty regarding the burden of influenza among children in some countries, the field effectiveness of inactivated and live attenuated vaccines when administered to children, the level of uptake achievable by national programmes and the amount of indirect protection that they will provide.
- There are also likely to be concerns regarding the resources required to expand

seasonal influenza vaccination to children in countries where programmes are publicly funded.

- Despite these concerns, there remains a persuasive argument for vaccinating children against influenza, especially in countries with the existing infrastructure to deliver such programmes but where there has been suboptimal uptake among at risk groups with their current programmes.
- As more data become available on the effectiveness of vaccinating European children against influenza, for example through the UK programme, European countries are likely to be faced with the decision whether to continue the traditional approach of targeting groups at the highest risk of complications (and for whom the response to the vaccine is potentially poorest) or focus more on those most responsible for onward transmission.

### **Financial and competing interests disclosure**

A McGuire and M Drummond received an educational grant from AstraZeneca which allowed this review to be undertaken. The research undertaken by the London School of Economics was funded by AstraZeneca. The manuscript is based on a report produced by the London School of Economics, funded by AstraZeneca, available

at: <http://blogs.lse.ac.uk/lseenterprise/files/2015/04/ChildhoodAndAdolescentInfluenzaVaccination.pdf>. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from

those disclosed. Editorial support was provided by Talya Underwood, Prime Medical Group, London, UK and Luke Cripps, Ogilvy Healthworld, London, UK funded by AstraZeneca. The opinions, conclusions, and interpretation of the data are the responsibility of the authors.

## References

### *Reference annotations*

*\* Of interest*

*\*\* Of considerable interest*

1. World Health Organization. Influenza (Seasonal) Fact sheet N°211. 2014.  
Available at: <http://www.who.int/mediacentre/factsheets/fs211/en/>. [Last accessed 3 February 2015].
2. Rothberg MB, Haessler SD, Brown RB. Complications of viral influenza. *Am J Med* 2008;121:258-64.
3. European Centre for Disease Prevention and Control. Influenza remains a threat. 2015. Available  
at: [http://www.ecdc.europa.eu/en/healthtopics/seasonal\\_influenza/key\\_messages/Pages/key\\_messages\\_2.aspx](http://www.ecdc.europa.eu/en/healthtopics/seasonal_influenza/key_messages/Pages/key_messages_2.aspx). [Last accessed 3 February 2015].
4. Simonsen L, Fukuda K, Schonberger LB, et al. The impact of influenza epidemics on hospitalizations. *J Infect Dis* 2000;181:831-7.
5. Fowlkes A, Dasgupta S, Chao E, et al. Estimating influenza incidence and rates of influenza-like illness in the outpatient setting. *Influenza Other Respir Viruses* 2013;7:694-700.
6. Mossong J, Hens N, Jit M, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* 2008;5:e74.

7. Cromer D, van Hoek AJ, Jit M, et al. The burden of influenza in England by age and clinical risk group: a statistical analysis to inform vaccine policy. *J Infect* 2014;68:363-71.
8. Fraaij PL, Heikkinen T. Seasonal influenza: the burden of disease in children. *Vaccine* 2011;29:7524-8.
9. Heikkinen T, Silvennoinen H, Peltola V, et al. Burden of influenza in children in the community. *J Infect Dis* 2004;190:1369-73.
10. Reed C, Kallen AJ, Patton M, et al. Infection with community-onset *Staphylococcus aureus* and influenza virus in hospitalized children. *Pediatr Infect Dis J* 2009;28:572-6.
11. Toovey S. Influenza-associated central nervous system dysfunction: a literature review. *Travel Med Infect Dis* 2008;6:114-24.
12. Heikkinen T, Ruuskanen O, Waris M, et al. Influenza vaccination in the prevention of acute otitis media in children. *Am J Dis Child* 1991;145:445-8.
13. Block SL, Heikkinen T, Toback SL, et al. The efficacy of live attenuated influenza vaccine against influenza-associated acute otitis media in children. *Pediatr Infect Dis J* 2011;30:203-7.
14. Grijalva CG, Zhu Y, Williams DJ, et al. Association Between Hospitalization With Community-Acquired Laboratory-Confirmed Influenza Pneumonia and Prior Receipt of Influenza Vaccination. *JAMA* 2015;314:1488-97.

15. Chow MY, Yin JK, Heron L, et al. The impact of influenza-like illness in young children on their parents: a quality of life survey. *Qual Life Res* 2014;23:1651-60.
16. Poehling KA, Edwards KM, Weinberg GA, et al. The underrecognized burden of influenza in young children. *N Engl J Med* 2006;355:31-40.
17. O'Reilly FW, Stevens AB. Sickness absence due to influenza. *Occup Med (Lond)* 2002;52:265-9.
18. Akazawa M, Sindelar JL, Paltiel AD. Economic costs of influenza-related work absenteeism. *Value Health* 2003;6:107-15.
19. Palmer LA, Rousculp MD, Johnston SS, et al. Effect of influenza-like illness and other wintertime respiratory illnesses on worker productivity: The child and household influenza-illness and employee function (CHIEF) study. *Vaccine* 2010;28:5049-56.
20. Centers for Disease Control and Prevention. Prevention and control of influenza with vaccines: recommendations of the Advisory Committee on Immunization Practices (ACIP), 2010. *MMWR Morb Mort Wkly Rep* 2010;59:1-62.
21. Public Health Agency of Canada. Statement on Seasonal Influenza Vaccine for 2011–2012. *CCDR* 2011;37:1-55.
22. Public Health Agency of Canada. Statement on Seasonal Influenza Vaccine for 2012–2013. *CCDR* 2012;38:

23. \*\* Antonova EN, Rycroft CE, Ambrose CS, et al. Burden of paediatric influenza in Western Europe: a systematic review. BMC Public Health 2012;12:968.

***This systematic review provides evidence that the pediatric burden of influenza in Europe is substantial and has a significant direct impact on children and an indirect impact on their siblings and parents. The authors propose that the evidence identified in this review could aid European childhood immunization policy decisions.***

24. European Centre for Disease Prevention and Control. ECDC scientific advice on seasonal influenza vaccination of children and pregnant women. 2012. Available at: [http://ecdc.europa.eu/en/publications/Publications/20120905\\_GUI\\_HP\\_V\\_vaccine\\_update.pdf](http://ecdc.europa.eu/en/publications/Publications/20120905_GUI_HP_V_vaccine_update.pdf). [Last accessed 12 March 2015].
25. Rahmqvist M, Gjessing K, Faresjö T. Influenza-like illness: Young children, working parents and primary care take the strain. 33rd Annual Meeting of the European Society for Paediatric Infectious Diseases (ESPID 2015) <http://espid2015.kenes.com/Documents/ESPID%202015%20Abstracts.pdf>.
26. European Commission. Employment in Europe in 2010. 2010. Available at: <http://ec.europa.eu/social/home.jsp?langId=en>. [Last accessed 1 February 2015].
27. Keeling MJ, Danon L. Mathematical modelling of infectious diseases. Br Med Bull 2009;92:33-42.

28. Plans-Rubio P. The vaccination coverage required to establish herd immunity against influenza viruses. *Prev Med* 2012;55:72-7.
29. Lambert LC, Fauci AS. Influenza vaccines for the future. *N Engl J Med* 2010;363:2036-44.
30. Heikkinen T, Ikonen N, Ziegler T. Impact of influenza B lineage-level mismatch between trivalent seasonal influenza vaccines and circulating viruses, 1999-2012. *Clin Infect Dis* 2014;59:1519-24.
31. Ambrose CS, Levin MJ. The rationale for quadrivalent influenza vaccines. *Hum Vaccin Immunother* 2012;8:81-8.
32. Newall AT, Jit M, Beutels P. Economic evaluations of childhood influenza vaccination: a critical review. *Pharmacoeconomics* 2012;30:647-60.
33. Rhorer J, Ambrose CS, Dickinson S, et al. Efficacy of live attenuated influenza vaccine in children: A meta-analysis of nine randomized clinical trials. *Vaccine* 2009;27:1101-10.
34. Vesikari T, Knuf M, Wutzler P, et al. Oil-in-water emulsion adjuvant with influenza vaccine in young children. *N Engl J Med* 2011;365:1406-16.
35. Caspard H, Gaglani M, Clipper L, et al. Effectiveness of live attenuated influenza vaccine and inactivated influenza vaccine in children 2-17 years of age in 2013-2014 in the United States. *Vaccine* 2015;
36. Monto AS, Davenport FM, Napier JA, et al. Modification of an outbreak of influenza in Tecumseh, Michigan by vaccination of schoolchildren. *J Infect Dis* 1970;122:16-25.



37. Piedra PA, Gaglani MJ, Kozinetz CA, et al. Herd immunity in adults against influenza-related illnesses with use of the trivalent-live attenuated influenza vaccine (CAIV-T) in children. *Vaccine* 2005;23:1540-8.
38. Ghendon YZ, Kaira AN, Elshina GA. The effect of mass influenza immunization in children on the morbidity of the unvaccinated elderly. *Epidemiol Infect* 2006;134:71-8.
39. Reichert TA, Sugaya N, Fedson DS, et al. The Japanese experience with vaccinating schoolchildren against influenza. *N Engl J Med* 2001;344:889-96.
40. Hak E, Verheij TJ, Grobbee DE, et al. Confounding by indication in non-experimental evaluation of vaccine effectiveness: the example of prevention of influenza complications. *J Epidemiol Community Health* 2002;56:951-5.
41. \* Loeb M, Russell ML, Moss L, et al. Effect of influenza vaccination of children on infection rates in Hutterite communities: a randomized trial. *JAMA* 2010;303:943-50.

***This cluster randomized trial of children and adolescents aged 36 months to 15 years provides robust evidence of herd protection from influenza vaccination programs.***

42. \*King JC, Jr., Stoddard JJ, Gaglani MJ, et al. Effectiveness of school-based influenza vaccination. *N Engl J Med* 2006;355:2523-32.

***This intervention study of school-based influenza vaccination found that outcomes related to influenza-like illness were significantly lower in households from the intervention schools that received influenza***

***vaccination than in households from the control-school that did not receive influenza vaccination.***

43. Fiore AE, Uyeki TM, Broder K, et al. Prevention and control of influenza with vaccines: recommendations of the Advisory Committee on Immunization Practices (ACIP), 2010. MMWR Recomm Rep 2010;59:1-62.
44. European Centre for Disease Prevention and Control. Immunisation schedules by target disease. 2015. Available at: <http://vaccine-schedule.ecdc.europa.eu/Pages/Scheduler.aspx>. [Last accessed 1 February 2015].
45. Vaccine European New Integrated Collaborative Effort (VENICE). Final report on Seasonal Influenza Vaccination survey in EU/EEA countries, Season 2011-2012 (amended version). 2013. Available at: [http://venice.cineca.org/VENICE\\_Seasonal\\_Influenza\\_2011-12\\_1.2v.pdf](http://venice.cineca.org/VENICE_Seasonal_Influenza_2011-12_1.2v.pdf). [Last accessed 1 February 2015].
46. Walkinshaw E. Mandatory vaccinations: The international landscape. CMAJ 2011;183:E1167-E1168.
47. Mereckiene J, Cotter S, Nicoll A, et al. Seasonal influenza immunisation in Europe. Overview of recommendations and vaccination coverage for three seasons: pre-pandemic (2008/09), pandemic (2009/10) and post-pandemic (2010/11). Euro Surveill 2014;19:20780.
48. Kunze U, Bohm G, Groman E. Influenza vaccination in Austria from 1982 to 2011: a country resistant to influenza prevention and control. Vaccine 2013;31:5099-103.

49. Damm O, Eichner M, Rose MA, et al. Public health impact and cost-effectiveness of intranasal live attenuated influenza vaccination of children in Germany. *Eur J Health Econ* 2015;16:471-88.
50. Elonsalo, U. The Finnish Influenza Stakeholder Network: case report. 2009. Available at: <http://www.eswi.org/activities/country-influenza-stakeholder-networks/the-finnish-influenza-stakeholder-network-case-report>. [Last accessed 1 February 2015].
51. Blank PR, Schwenkglenks M, Szucs TD. Vaccination coverage rates in eleven European countries during two consecutive influenza seasons. *J Infect* 2009;58:446-58.
52. Joint Committee on Vaccination and Immunisation. Statement on the annual influenza vaccination programme – extension of the programme to children. 2012. Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/224775/JCVI-statement-on-the-annual-influenza-vaccination-programme-25-July-2012.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/224775/JCVI-statement-on-the-annual-influenza-vaccination-programme-25-July-2012.pdf). [Last accessed 10 March 2015].
53. Pebody RG, Green HK, Andrews N, et al. Uptake and impact of a new live attenuated influenza vaccine programme in England: early results of a pilot in primary school-age children, 2013/14 influenza season. *Euro Surveill* 2014;19:
54. \*\*Pebody RG, Green HK, Andrews N, et al. Uptake and impact of vaccinating school age children against influenza during a season with circulation of drifted influenza A and B strains, England, 2014/15. *Euro Surveill* 2015;20:

***This publication shares the uptake and impact of the UK's childhood influenza vaccination program during its second season in 2014–2015, demonstrating that vaccinating primary school age children in pilot areas resulted in significant and non-significant reductions in a range of surveillance indicators for both the targeted age groups and wider society (due to herd protection).***

55. European Medicines Agency. Guideline on the clinical evaluation of new vaccines. 2006. Available at: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2009/09/WC500003870.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003870.pdf). [Last accessed 1 February 2015].
56. Ambrose CS, Wu X, Knuf M, et al. The efficacy of intranasal live attenuated influenza vaccine in children 2 through 17 years of age: a meta-analysis of 8 randomized controlled studies. *Vaccine* 2012;30:886-92.
57. Osterholm MT, Kelley NS, Sommer A, et al. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *Lancet Infect Dis* 2012;12:36-44.
58. Carmona A, Omenaca F, Tejedor JC, et al. Immunogenicity and safety of AS03-adjuvanted 2009 influenza A H1N1 vaccine in children 6-35 months. *Vaccine* 2010;28:5837-44.
59. European Medicines Agency. Withdrawal Assessment Report Fluad paediatric. 2012. Available at: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Application\\_with](http://www.ema.europa.eu/docs/en_GB/document_library/Application_with)

[drawal assessment report/2012/04/WC500126030.pdf](http://ecdc.europa.eu/en/publications/Publications/Vaesco%20report%20Final%20with%20cover.pdf). [Last accessed 5 August 2015].

60. European Centre for Disease Prevention and Control. Narcolepsy in association with pandemic influenza vaccination. 2012. Available at: <http://ecdc.europa.eu/en/publications/Publications/Vaesco%20report%20Final%20with%20cover.pdf>. [Last accessed 7 August 2015].
61. Heinonen S, Silvennoinen H, Lehtinen P, et al. Effectiveness of inactivated influenza vaccine in children aged 9 months to 3 years: an observational cohort study. *Lancet Infect Dis* 2011;11:23-9.
62. Kissling E, Valenciano M, Buchholz U, et al. Influenza vaccine effectiveness estimates in Europe in a season with three influenza type/subtypes circulating: the I-MOVE multicentre case-control study, influenza season 2012/13. *Euro Surveill* 2014;19:
63. Pebody R, Warburton F, Andrews N, et al. Effectiveness of seasonal influenza vaccine in preventing laboratory-confirmed influenza in primary care in the United Kingdom: 2014/15 end of season results. *Euro Surveill* 2015;20:
64. Heikkinen T, Tsolia M, Finn A. Vaccination of healthy children against seasonal influenza: a European perspective. *Pediatr Infect Dis J* 2013;32:881-8.
65. Jit M, Newall AT, Beutels P. Key issues for estimating the impact and cost-effectiveness of seasonal influenza vaccination strategies. *Hum Vaccin Immunother* 2013;9:834-40.

66. Trotter CL, Edmunds WJ. Modelling cost effectiveness of meningococcal serogroup C conjugate vaccination campaign in England and Wales. *BMJ* 2002;324:809.

67. \*Pitman RJ, White LJ, Sculpher M. Estimating the clinical impact of introducing paediatric influenza vaccination in England and Wales. *Vaccine* 2012;30:1208-24.

***This assessment of the population impact of implementing a programme of pediatric vaccination finds that vaccinating as few as 50% of 2–18 year olds could significantly reduce annual incidence of influenza-related morbidity and mortality across the population and offer herd protection to the wider community.***

68. \*Baguelin M, Flasche S, Camacho A, et al. Assessing optimal target populations for influenza vaccination programmes: an evidence synthesis and modelling study. *PLoS Med* 2013;10:e1001527.

***This study proposes a framework for the integration of influenza surveillance data into transmission models to assess the optimal target populations for influenza vaccination programs. The model finds that the most efficient way to reduce overall influenza-attributable morbidity and mortality is to target children; consequently, the authors suggest that high-income countries should consider revising their influenza vaccination programs.***

69. Salo H, Kilpi T, Sintonen H, et al. Cost-effectiveness of influenza vaccination of healthy children. *Vaccine* 2006;24:4934-41.

70. Schmitt HJ, Booy R, Weil-Olivier C, et al. Child vaccination policies in Europe: a report from the Summits of Independent European Vaccination Experts. *Lancet Infect Dis* 2003;3:103-8.
71. Humiston SG, Rosenthal SL. Challenges to vaccinating adolescents: vaccine implementation issues. *Pediatr Infect Dis J* 2005;24:S134-S140.
72. European Centre for Disease Prevention and Control. Current practices in immunisation policymaking in European Countries. 2015. Available at: <http://www.ecdc.europa.eu/en/publications/Publications/Current-practices-on-immunisation-policymaking-processes-Mar-2015.pdf>. [Last accessed 10 March 2015].
73. Ricciardi GW, Toumi M, Weil-Olivier C, et al. Comparison of NITAG policies and working processes in selected developed countries. *Vaccine* 2015;33:3-11.
74. Joint Committee on Vaccination and Immunisation. Code of Practice. 2013. Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/224864/JCVI Code of Practice revision 2013 - final.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/224864/JCVI_Code_of_Practice_revision_2013_-_final.pdf). [Last accessed 10 March 2015].
75. NHS England. A modern data service for the NHS. 2013. Available at: <http://www.england.nhs.uk/ourwork/tsd/care-data/better-care/>. [Last accessed 10 March 2015].
76. Danzon PM, Sousa Pereira N. Vaccine Supply: Effects of Regulation and Competition. *Int J Econ Bus* 2011;18:239-71.

77. Department of Health. Getting it right for children, young people and families. Maximising the contribution of the school nursing team: Vision and call to action. 2012. Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/216464/dh\\_133352.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216464/dh_133352.pdf). [Last accessed 10 March 2015].
78. Olivier CW. Influenza vaccination coverage rate in children: reasons for a failure and how to go forward. Hum Vaccin Immunother 2012;8:107-18.
79. Cooper Robbins SC, Ward K, Skinner SR. School-based vaccination: a systematic review of process evaluations. Vaccine 2011;29:9588-99.
80. \*\*Kassianos G, White S, Reynolds AJ, et al. Review of the experiences from the first childhood influenza vaccination programme with a live attenuated influenza vaccine in England and Scotland. Drugs Context 2015;4:212280.

***This article reviews the experience from the first year of the UK's childhood influenza vaccination program and describes the processes utilized in England and Scotland for program setup, workforce management, identification and care of contraindicated patients, collection of data on vaccine uptake, communication strategies, and education of parents and children. This article shares lessons learnt to provide guidance for other countries considering the implementation of childhood influenza vaccination programs.***

81. Kwong JC, Pereira JA, Quach S, et al. Randomized evaluation of live attenuated vs. inactivated influenza vaccines in schools (RELATIVES) pilot study: a cluster randomized trial. Vaccine 2015;33:535-41.



82. Centers for Disease Control and Prevention. National Early Season Flu Vaccination Coverage, United States, November 2014. 2014. Available at: <http://www.cdc.gov/flu/fluview/nifs-estimates-nov2014.htm>. [Last accessed 5 August 215].
83. Briss PA, Rodewald LE, Hinman AR, et al. Reviews of evidence regarding interventions to improve vaccination coverage in children, adolescents, and adults. The Task Force on Community Preventive Services. *Am J Prev Med* 2000;18:97-140.
84. Blank P, Schwenkglens M, Szucs TD. The impact of European vaccination policies on seasonal influenza vaccination coverage rates in the elderly. *Hum Vaccin Immunother* 2012;8:328-35.
85. Yaqub O, Castle-Clarke S, Sevdalis N, et al. Attitudes to vaccination: a critical review. *Soc Sci Med* 2014;112:1-11.
86. Flodgren G, Eccles MP, Shepperd S, et al. An overview of reviews evaluating the effectiveness of financial incentives in changing healthcare professional behaviours and patient outcomes. *Cochrane Database Syst Rev* 2011;CD009255.
87. Bansal S, Pourbohloul B, Hupert N, et al. The shifting demographic landscape of influenza. *PLoS Curr* 2009;1:RRN1047.
88. Cipollone A, Patacchini E, Vallanti G. Female labour market participation in Europe: novel evidence on trends and shaping factors. *IZA Journal of European Labor Studies* 2014;3:18.

89. GOV.UK Letter. Vaccines and gelatine: PHE response. 2014. Available at: <https://www.gov.uk/government/news/vaccines-and-gelatine-phe-response>. [Last accessed 25 June 2015].
90. Leask J, Chapman S, Hawe P, et al. What maintains parental support for vaccination when challenged by anti-vaccination messages? A qualitative study. *Vaccine* 2006;24:7238-45.
91. Goldacre B. Medicine and the media: MMR: the scare stories are back. *BMJ* 2007;335:126-7.
92. Gupta RK, Best J, MacMahon E. Mumps and the UK epidemic 2005. *BMJ* 2005;330:1132-5.
93. Childhood flu immunisation programme: update and provisional roll-out schedule. 2014. Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/331045/2014-15\\_update\\_and\\_roll-out\\_schedule\\_letter.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/331045/2014-15_update_and_roll-out_schedule_letter.pdf). [Last accessed 11 November 2014].
94. European Centre for Disease Prevention and Control. Introduction of HPV vaccines in European Union countries – an update. 2012. Available at: [http://ecdc.europa.eu/en/publications/Publications/20120905\\_GUI\\_HPV\\_vaccine\\_update.pdf](http://ecdc.europa.eu/en/publications/Publications/20120905_GUI_HPV_vaccine_update.pdf). [Last accessed 12 March 2015].

ACCEPTED MANUSCRIPT

**Table 1. Regions and their constituent countries**

Regions					
Baltic	Central	Eastern	Nordic	Southern	Western
Estonia	Austria	Bulgaria	Denmark	Cyprus	Belgium
Latvia	Germany	Croatia	Finland	Greece	France
Lithuania	Liechtenstein	Czech R.	Iceland	Italy	Ireland
	Poland	Hungary	Norway	Malta	Luxembourg
		Romania	Sweden	Portugal	Netherlands
		Slovakia		Spain	United Kingdom
		Slovenia			


**Table 2. Overview of health system and vaccine policy making features for a sample of European countries**

Country	Main system of finance	Final decision maker on immunization	Funding source for immunization schedule	Tender system in place	Main provider of immunizations	NITAG	Consideration of cost-effectiveness required
Austria	Compulsory social insurance	Ministry of Health after negotiations with other stakeholders	Mixed	National level	Private practice	Yes	Yes
Belgium	Compulsory social insurance	National and regional Ministries of Health	Mixed (childhood vaccines tax-funded)	Regional level	Private practice	Yes	No
Denmark	Taxation	Parliament	Tax-funded	National level	Public providers	Yes	Yes
Finland	Taxation	Parliament with recommendation from Ministry of Finance	Tax-funded	National level	Public providers	Yes	Yes
France	Compulsory social insurance	Ministry of Health	Mixed	National level	Private practice	Yes	Yes
Germany	Compulsory social insurance	NITAG	Social insurance	Regional level (for some vaccines)	Private practice	Yes	Yes
Greece	Taxation	Ministry of Health	Tax-funded	National level	Private practice	Yes	No
Ireland	Taxation	Ministry of Health	Tax-funded	National level	Public providers, including school nurses	Yes	Yes
Italy	Taxation	National and regional Ministries of Health	Tax-funded	National, regional and local level	Public providers	Yes	Yes
Netherlands	Mixed	Ministry of Health	Tax-funded	National level	Public providers	Yes	Yes
Portugal	Taxation	Ministry of Health	Tax-funded	No system in place	Public providers	Yes	Yes
Spain	Taxation	National and regional Ministries of Health	Tax-funded	No system in place	Public providers, including school nurses	Yes	Yes
Sweden	Taxation	Government	Tax-funded	Regional and local level	Public providers, including school nurses	No	Yes
UK	Taxation	Ministry of Health	Tax-funded	National level	Public providers, including school nurses	Yes	Yes


Data on whether or not school-based immunization takes place were taken from a review of human papillomavirus vaccination programmes [94]

NITAG: National Immunisation Technical Advisory Groups


**Figure 1. Childhood and adolescent age groups with a general recommendation for seasonal influenza vaccination in Europe**



**Figure 2. Number of countries with a general recommendation for the vaccination of children and adolescents by geographical region**



**Figure 3. European countries ranked by the proportion of their population aged between 0 and 14 years in 2012**



Countries with a recommendation for healthy children to be vaccinated against influenza are in red, with the dashed line representing the median for all countries